



The Fat Trap™

For easy disposal of all kitchen fats & oils

Every year, thousands of litres of hot fat and cooking oil are poured down kitchen sinks. When fat and oil cool they build up on the pipes and can cause a blockage in your drains, leading to inconvenience, and an expensive plumber's bill for the homeowner.

You can avoid this problem by collecting all your kitchen fats and oils in a **Fat Trap™**.

Once full, soften any fat by floating the **Fat Trap (lid on!)** for a few minutes in the hot water while you do the washing-up. Then pour the now-liquid contents into a throwaway container with a screw top. Put that in with the household rubbish.

To find out how to obtain a **Fat Trap**, contact **FOG-Catcher**.

Website: www.fattrap.co.nz


Phone: **0800 FAT TRAP (32 88 72)**

Write to: **FOG-Catcher Limited, PO Box 35210,
Naenae, Lower Hutt 5041**

Keep **your** drains free of blockages
- with a **Fat Trap**

Use a **Fat Trap** to help keep the environment clean and prevent blocked drains

Clean, convenient and easy to use



The **Fat Trap** is a re-usable container for collecting waste fats and oils.

Did you know?...

A typical call-out charge to unblock a sink is \$150.

Make your very own "Bird Cake"

The fat content of your **Fat Trap** can be mixed with other things to make "Bird Cake", nutritious food for birds when food is scarce in the winter. Mixing it all up is messy but it is very helpful in Winter weather.

Drain off any oil, heat fat briefly in a pan to soften it, then mix in all sorts of seeds. Roll into a ball with a knotted string through it.



<http://nzbirds.com/more/feed.html> has lots of NZ-relevant information.

Advice from the Vet

It is unwise to supply bird cake during warm weather as the fat quickly goes rancid.

It is unwise to supply bird cake in the Spring as birds should be seeking out higher protein foods for reproduction and chick rearing and they may take the easy way out and become weakened.

Avoid feeding birds anything containing dairy products, excessively salty foods such as salted peanuts, or avocados (extremely toxic to birds). Do not give desiccated coconut or any dehydrated food as it will swell inside the bird. Meat scraps, most other fruits and veggies are fine, provided they are fresh.

Many birds of many types visit feeding stations, so they are ideal places for spreading diseases and fleas. Keep feeding stations and any water/nectar sources or bird baths scrupulously clean, and place them in areas out of reach of cats, etc.

Remove bird feeders if there is a bird-flu or similar outbreak so as to prevent it from spreading.